

On curvature homogeneous 4D Lorentzian manifolds

R. Milson, N. Pelavas

Dept. Mathematics and Statistics, Dalhousie University
Halifax NS B3H 3J5, Canada

E-mail: rmilson@dal.ca, pelavas@mathstat.dal.ca

Abstract. We prove that a four-dimensional Lorentzian manifold that is curvature homogeneous of order 3, or CH_3 for short, is necessarily locally homogeneous. We also exhibit and classify four-dimensional Lorentzian, CH_2 manifolds that are not homogeneous.

PACS numbers: 04.20, 02.40

AMS classification scheme numbers: 53C50

1. Introduction

A pseudo-Riemannian manifold (M, g_{ij}) is locally homogeneous if the Lie algebra of Killing vector fields spans the tangent space at all points of M . We say that M is curvature homogeneous of order s , or equivalently that M belongs to class CH_s if the components of the curvature tensor and its first s covariant derivatives are constant relative to some local frame.

Local homogeneity implies curvature homogeneity of all orders. Remarkably, the converse is also true, in some fashion. Let us say that a manifold is *properly* CH_s if it belongs to class CH_s , but is not locally homogeneous. Singer showed that there does not exist a proper CH_s manifold if s is larger than a certain invariant, called the Singer index [16, 19]. More precisely, we have the following.

Theorem 1.1 (Singer's criterion) *Let (M, g_{ij}) be a CH_s manifold of signature (p, q) . Let $\mathfrak{g}_r \subseteq \mathfrak{g}_{-1} := \mathfrak{so}(p, q)$, $r = 0, 1, \dots, s$ be the subalgebra that leaves invariant the arrays of constants $R^a_{b_1 b_2 b_3}, \nabla_c R^a_{b_1 b_2 b_3}, \dots, \nabla_{c_1 \dots c_r} R^a_{b_1 b_2 b_3}$. If $\mathfrak{g}_{s-1} = \mathfrak{g}_s$, then M is locally homogeneous.*

Many examples of proper CH_0 Riemannian [14] and Lorentzian manifolds [1] are known. In the case of three-dimensional curvature homogeneous geometries, classification results are available [3, 15]. See [10] for additional references, and for examples of higher-dimensional proper CH manifolds of general signature. In 4 dimensions, there are no proper CH_1 Riemannian manifolds[18], but this is not true in the Lorentzian[2] and the neutral-signature setting[5]. At this point, no examples of four-dimensional, proper CH_2 geometries seem to be known[11], leaving open the question of the value of the Singer index for four-dimensional, Lorentzian, curvature-homogeneous manifolds.

In this note we show that there are no four-dimensional, proper CH₃ Lorentzian manifolds; see section 3. We also exhibit and classify all four-dimensional, proper CH₂ Lorentzian manifolds; see eqns. (24)-(26).

Our approach is to formulate the field equations for a CH geometry as an exterior differential system (EDS) on the second-order frame bundle[7]. The integrability constraints manifest as algebraic constraints (torsion in EDS parlance) on the curvature scalars. The classification is accomplished by deriving torsion-free configurations for the CH field equations corresponding to various algebraic types of the curvature tensor, and by using Singer's criterion to rule out the homogeneous subcases. See [12] for an introduction to EDS theory, see [6, 13] for a discussion of higher-order frames. See [8, 4] for some recent applications of EDS theory to relativity. We will use this method to classify proper CH₁ four-dimensional, Lorentz geometries in a forthcoming publication.

2. The EDS for curvature-homogeneous geometries

Let M be an n -dimensional, analytic manifold, and let g_{ab} be a fixed, non-degenerate bilinear form. Henceforth, we use g_{ab} to lower frame indices, which we denote by $a, b, c = 1, \dots, n$. We use u^i to denote local coordinates, where $i, j = 1, \dots, n$ are coordinate indices. Even though we are interested in the case of $n = 4$ and g_{ab} of Lorentzian signature, much of the underlying theory can be given without these assumptions.

Pseudo-Riemannian structure can be specified in two ways: as a metric, or as a g_{ab} -orthogonal moving frame. In the second approach, instead of using g_{ij} and Γ^i_{jk} as unknowns in the field equations, the field variables are scalars

$$u^i_a, \det(u^i_a) \neq 0, \quad u^i_{bc} = u^i_{cb},$$

the $\frac{1}{2}n^2(n+3)$ components of a 2nd order moving frame [13]. The metric and connection can then be recovered as

$$\begin{aligned} g_{ij} &= v^a_i v^b_j g_{ab}, \\ \Gamma^i_{jk} &= -v^b_j v^c_k u^i_{bc}, \end{aligned}$$

where $v^a_i u^i_b = \delta^a_b$ is the inverse matrix of coframe components. It will be convenient to formulate the field equations in terms of 1-form variables

$$\omega^a = v^a_i du^i, \tag{1}$$

$$\omega^a_b = v^a_i (du^i_b - u^i_{bc} \omega^c), \tag{2}$$

which represent the coframe and connection 1-forms. The equations

$$d\omega^a + \omega^a_b \wedge \omega^b = 0, \tag{3}$$

$$(d\omega^a_b + \omega^a_c \wedge \omega^c_b) \wedge \omega^b = 0, \tag{4}$$

the 1st structure equations and algebraic Bianchi relations, follow as tautologies.

Let $G = \{X^a_b : X_{ca}X^c_b = g_{ab}\}$ and $\mathfrak{g} = \{M^a_b : M_{(ab)} = 0\}$ denote, respectively, the $N := \frac{1}{2}n(n-1)$ -dimensional group of g_{ab} -orthogonal transformations and the corresponding Lie algebra of anti-self dual infinitesimal transformations. Let $M_\alpha = (M^a_{b\alpha})$ be a basis of $\mathfrak{g}$, and let $C^\alpha_{\beta\gamma}$ denote the corresponding structure constants. Henceforth, we use bivector indices $\alpha, \beta, \gamma = 1, \dots, N$ as gauge indices. Let Γ^α denote the $\mathfrak{g}$ -valued portion of the connection 1-forms; to be more precise, $\omega_{[ab]} = M_{ab\alpha} \Gamma^\alpha$.

Particular instances of a CH_0 geometry are determined by an array of constants $\tilde{R}_{bc}^\alpha = -\tilde{R}_{cb}^\alpha$. The corresponding CH_0 field equations are as follows:

$$\omega_{(ab)} = 0, \quad (5)$$

$$d\omega_{(ab)} = 0, \quad (6)$$

$$d\Gamma^\alpha + \frac{1}{2}C_{\beta\gamma}^\alpha \Gamma^\beta \wedge \Gamma^\gamma - \frac{1}{2}\tilde{R}_{bc}^\alpha \omega^b \wedge \omega^c = 0, \quad (7)$$

$$(\mathbf{M}_\beta \cdot \tilde{R})_{bc}^\alpha \Gamma^\beta \wedge \omega^b \wedge \omega^c = 0, \quad (8)$$

where

$$(\mathbf{M}_\beta \cdot \tilde{R})_{bc}^\alpha = C_{\beta\gamma}^\alpha \tilde{R}_{bc}^\gamma - M_{b\beta}^a \tilde{R}_{ac}^\alpha - M_{c\beta}^a \tilde{R}_{ba}^\alpha \quad (9)$$

is the standard action of $\mathfrak{g}$ on the vector space of $\mathfrak{g}$ -valued curvature-type tensors. Equations (5) constrain the connection to be $\mathfrak{g}$ -valued; this is equivalent to the compatibility of Γ_{jk}^i and g_{ij} . Equations (6) and (8) are the compatibility conditions for (5) and (7), respectively; one takes the exterior derivative of the latter to obtain the former. In particular, equations (6) constrain the curvature 2-form,

$$\Omega_b^a := d\omega_b^a + \omega_c^a \wedge \omega_b^c,$$

to be $\mathfrak{g}$ -valued. Equations (8) express the differential Bianchi identities. Equations (7) (these are equivalent to the NP equations in 4D relativity) together with (6) express the condition for curvature homogeneity, namely

$$\Omega_b^a = \frac{1}{2}M_{b\alpha}^a \tilde{R}_{cd}^\alpha \omega^c \wedge \omega^d.$$

Hence, as a consequence of (4), it is necessary to assume that the $\tilde{R}_{bc}^\alpha$ satisfy the algebraic Bianchi identity, namely

$$\tilde{R}_{[bc}^\alpha M_{d]\alpha}^a = 0. \quad (10)$$

2.1. The CH_0 class

Set $G_{-1} := G$ and $\mathfrak{g}_{-1} := \mathfrak{g}$, and let $G_0 \subseteq G_{-1}$ and $\mathfrak{g}_0 \subseteq \mathfrak{g}_{-1}$ be, respectively, the subgroup and the Lie subalgebra of frame transformations that leaves invariant the constant array $\tilde{R}_{bc}^\alpha$. As a system, equations (5)-(8) are invariant with respect to G_0 . Thus, G_0 serves as the reduced gauge group of the CH_0 field equations. Let us set $N_i = \dim \mathfrak{g}_i$ and arrange the basis of $\mathfrak{g}$ so that $\mathbf{M}_\lambda$, $\lambda = 1, \dots, N_{-1} - N_0$ is a basis of $\mathfrak{g}_{-1}/\mathfrak{g}_0$ and so that $\mathbf{M}_\xi$, $\xi = N_{-1} - N_0 + 1, \dots, N_{-1}$ is a basis of the reduced subalgebra $\mathfrak{g}_0$. It is important to note that if $\mathfrak{g}_0 = \mathfrak{g}_{-1}$, then the Bianchi identities (8) reduce to a trivial constraint. Consequently, equations (3) and (7) function as the structure equations of a homogeneous space — in this case a space of constant curvature. Thus, to obtain a proper CH_0 geometry it is necessary to assume that the inclusion $\mathfrak{g}_0 \subset \mathfrak{g}_{-1}$ is proper.

The components of the connection forms Γ^α do not obey a meaningful transformation law with respect to the full group G . However, once we restrict our group to G_0 , we distinguish between two different kinds of connection forms : Γ^ξ , whose components are not G_0 -equivariant, and

$$\Gamma^\lambda = \Gamma_a^\lambda \omega^a, \quad (11)$$

whose G_0 -equivariant components Γ_a^λ we treat as new scalar variables. By (2), these new variables represent certain linear combinations of the first order jets

$$u_{a;b}^i = \partial_j u_a^i u_b^j - u_{ab}^i.$$

The Γ_a^λ obey the following field equations:

$$d\Gamma_a^\lambda \wedge \omega^a + (M_\xi \cdot \Gamma)_a^\lambda \Gamma^\xi \wedge \omega^a + \frac{1}{2} \Upsilon_{bc}^\lambda \omega^b \wedge \omega^c = 0, \quad (12)$$

where

$$\Upsilon_{bc}^\lambda = -2\Gamma_a^\lambda \Gamma_{[b}^\mu M_{c]\mu}^a + \Gamma_b^\mu \Gamma_c^\nu C_{\mu\nu}^\lambda - \tilde{R}_{bc}^\lambda. \quad (13)$$

Here

$$(M_\xi \cdot \Gamma)_a^\lambda = C_{\xi\mu}^\lambda \Gamma_b^\mu - M_{a\xi}^b \Gamma_b^\lambda \quad (14)$$

denotes the natural action of $\mathfrak{g}_0$ on $\mathcal{L}(\mathbb{R}^n, \mathfrak{g}_{-1}/\mathfrak{g}_0)$, the space of linear maps from $\mathbb{R}^n$ to $\mathfrak{g}_{-1}/\mathfrak{g}_0$. Eqns. (12) follow from (7) and from the exterior derivative of (11).

Taken together, (5)-(8) generate an exterior differential system. The total space is $\mathcal{F}^2 M$, the bundle of 2nd order frames over M [13]. The variables u^i, u_a^i, u_{bc}^i serve as local coordinates on $\mathcal{F}^2 M$. We impose the independence condition

$$\omega^1 \wedge \cdots \wedge \omega^n \neq 0, \quad (15)$$

which means that u^i are to be regarded as independent variables, and u_a^i, u_{bc}^i as the field variables. An n -dimensional integral manifold with independence condition (15) describes a g_{ab} -orthogonal moving frame with constant R_{bc}^α . From a theoretical point of view, however, it is more advantageous to consider $(n + N_0)$ -dimensional integral manifolds with independence condition

$$\left(\bigwedge_a \omega^a \right) \wedge \left(\bigwedge_\xi \Gamma^\xi \right) \neq 0. \quad (16)$$

The latter type of integral manifold describes a CH_0 bundle of g_{ab} -orthogonal frames with reduced structure group G_0 .

For general choices of the $\tilde{R}_{bc}^\alpha$ the EDS is not involutive. We must (partially) prolong the system by introducing the extra field variables Γ_a^λ ; the prolonged total space is $\mathcal{F}^2 M \times \mathcal{L}(\mathbb{R}^n, \mathfrak{g}_{-1}/\mathfrak{g}_0)$. The prolonged EDS is generated by equations (5)-(8), (11), (12). Using (11), the Bianchi identities take the form of linear constraints

$$(M_\lambda \cdot \tilde{R})_{[bc}^\alpha \Gamma_{a]}^\lambda = 0. \quad (17)$$

Some linear combinations of (12) reduce to algebraically quadratic constraints on the Γ_a^λ . In EDS theory, such algebraic constraints are called torsion [12]. In order for the prolonged equations to have solutions, it is necessary to constrain the prolongation variables so that all torsion vanishes.

2.2. Higher order curvature homogeneity

On a manifold of class CH_1 , the curvature tensor R_{bc}^α and its covariant derivative $\nabla_a R_{bc}^\alpha$ are constant relative to some choice of moving frame. Consequently, there is a chain of 3 Lie algebras, call them $\mathfrak{g}_1 \subseteq \mathfrak{g}_0 \subseteq \mathfrak{g}_{-1}$, such that $R_{bc}^\alpha = \tilde{R}_{bc}^\alpha$ and $\nabla_a R_{bc}^\alpha = \tilde{R}_{abc}^\alpha$, where the latter are respectively, a $\mathfrak{g}_0$ -invariant and a $\mathfrak{g}_1$ -invariant array of constants. We also have 3 groups of generators: M_ρ a basis of $\mathfrak{g}_{-1}/\mathfrak{g}_0$, M_λ a basis of $\mathfrak{g}_0/\mathfrak{g}_1$, and M_ξ a basis of $\mathfrak{g}_1$. In other words, a choice of $\tilde{R}_{bc}^\alpha$ reduces the gauge/structure group from $\mathfrak{g}_{-1}$ to $\mathfrak{g}_0$; then, $\tilde{R}_{abc}^\alpha$ further reduces the group to $\mathfrak{g}_1$.

Since the $\tilde{R}_{bc}^\alpha$ are constant, we have

$$\tilde{R}_{abc}^\alpha = (M_\rho \cdot \tilde{R})_{bc}^\alpha \Gamma_a^\rho. \quad (18)$$

Since the M_ξ, M_λ span the kernel of the action of $\mathfrak{g}_{-1}$ on $\tilde{R}_{bc}^\alpha$, by definition, one can fully back-solve the linear system (18) to obtain Γ_a^ρ as a bilinear combination of $\tilde{R}_{bc}^\alpha$ and $\tilde{R}_{abc}^\alpha$. In describing a CH_1 geometry, it therefore suffices to specify a $\mathfrak{g}_1$ -invariant array of constants, $\Gamma_a^\rho = \tilde{\Gamma}_a^\rho$, rather than an array of constants $\tilde{R}_{abc}^\alpha$.

The EDS for CH_1 geometry is given by equations (5)-(8), and by

$$\Gamma^\rho - \tilde{\Gamma}_a^\rho \omega^a = 0 \quad (19)$$

$$(M_\lambda \cdot \tilde{\Gamma})_a^\rho \Gamma^\lambda \wedge \omega^a + \frac{1}{2} \tilde{\Upsilon}_{bc}^\rho \omega^b \wedge \omega^c = 0, \quad (20)$$

where $\tilde{\Upsilon}_{bc}^\rho$ are constants that are given, mutatis mutandi, by (13). The Bianchi equations (17) act as linear constraints on the $\tilde{\Gamma}_a^\rho$. Certain linear combinations of (20) may act as quadratic constraints. Even if these constraints are satisfied, the resulting EDS is, in general, not involutive; we have to prolong. As above, we introduce additional scalar variables Γ_a^λ and additional field equations (11) and (12). As a consequence of (11) and (20), we have the linear constraints

$$(M_\lambda \cdot \tilde{\Gamma})_{[b}^\rho \Gamma_{c]}^\lambda = \frac{1}{2} \tilde{\Upsilon}_{bc}^\rho. \quad (21)$$

As well, some linear combinations of the field equations (12) reduce to algebraically quadratic constraints on the Γ_a^λ . Thus, it is necessary to constrain the $\tilde{\Gamma}_a^\rho$ and Γ_a^λ so that all torsion implied by (17), (12), (20), (21) vanishes. We define a CH_1 configuration to be a triple $(\tilde{R}_{bc}^\alpha, \tilde{\Gamma}_a^\rho, V_1)$, where the $\tilde{R}_{bc}^\alpha$ is an array of $\mathfrak{g}_0$ -invariant constants that satisfies (10), where $\tilde{\Gamma}_a^\rho$ is an array of $\mathfrak{g}_1$ -invariant constants, and where $V_1 \subseteq \mathcal{L}(\mathbb{R}^n, \mathfrak{g}_0/\mathfrak{g}_1)$ is an algebraic variety such that the EDS generated by (5)-(8), (11), (12), (19), (20) on $\mathcal{F}^2 M \times V_1$ is torsion-free.

An important subcase arises when $\mathfrak{g}_1 = \mathfrak{g}_0$ [9]. In this case, the constants $\tilde{\Gamma}_a^\rho$ must be $\mathfrak{g}_0$ -invariant, and must satisfy the linear and quadratic constraints

$$(M_\rho \cdot R)_{[bc}^\alpha \tilde{\Gamma}_{a]}^\rho = 0, \quad \tilde{\Upsilon}_{bc}^\rho = 0.$$

Since $\mathfrak{g}_0/\mathfrak{g}_1$ is trivial, no additional constraints can or need be imposed. The unprolonged EDS on $\mathcal{F}^2 M$, generated by (5)-(8), (19) is involutive. The $(n + N_0)$ -dimensional integral manifolds of this EDS are homogeneous spaces with $\mathfrak{g}_0$ isotropy and structure equations

$$d\omega^a + M_{c\rho}^a \tilde{\Gamma}_b^\rho \omega^b \wedge \omega^c + M_{b\xi}^a \Gamma^\xi \wedge \omega^b = 0 \quad (22)$$

$$d\Gamma^\xi + \frac{1}{2} C_{v\zeta}^\xi \Gamma^v \wedge \Gamma^\zeta - \frac{1}{2} \tilde{R}_{bc}^\xi \omega^b \wedge \omega^c = 0. \quad (23)$$

We say that a CH_1 configuration is proper if the inclusion $\mathfrak{g}_1 \subset \mathfrak{g}_0$ is proper. In seeking proper CH_1 geometries one must demand that the CH_1 configuration be proper.

The general treatment of CH_s , $s \geq 1$ is similar. Now one has to consider a chain of Lie algebras $\mathfrak{g}_s \subseteq \dots \subseteq \mathfrak{g}_0 \subseteq \mathfrak{g}_{-1}$, with M_{ρ_r} , $r = 1, \dots, s$ a basis of $\mathfrak{g}_{r-2}/\mathfrak{g}_{r-1}$, with M_λ a basis for $\mathfrak{g}_{s-1}/\mathfrak{g}_s$, and M_ξ a basis for $\mathfrak{g}_s$. The linear coordinates, Γ_a^λ , on $\mathcal{L}(\mathbb{R}^n, \mathfrak{g}_{s-1}/\mathfrak{g}_s)$, are prolongation variables. A CH_s configuration is an $(s+2)$ -tuple $(\tilde{R}_{bc}^\alpha, \tilde{\Gamma}_a^{\rho_r}, V_s)$, where $\tilde{R}_{bc}^\alpha$ is an array of $\mathfrak{g}_0$ -invariant constants, where each $\tilde{\Gamma}_a^{\rho_r}$ is an array of $\mathfrak{g}_r$ -invariant constants, and where $V_s \subseteq \mathcal{L}(\mathbb{R}^n, \mathfrak{g}_{s-1}/\mathfrak{g}_s)$ is an algebraic variety such that the EDS generated by (5)-(8), (11), (12), (19), (20) on $\mathcal{F}^2 M \times V_s$ is torsion-free. The definition of involutive configuration is as above. An n -dimensional integral manifold with independence condition (15) describes a g_{ab} -orthogonal frame with constant $\nabla_{c_1 \dots c_r} R_{b_1 b_2 b_3}^\alpha$, $r = 0, 1, \dots, s$. An $(n + N_s)$ -dimensional integral manifold with independence condition (16) describes a bundle of g_{ab} -orthogonal frames with reduced structure group G_s .

3. Proper CH₂ geometries

Henceforth, we assume that M is four-dimensional and that g_{ab} is the Lorentzian inner product. We will express our calculations using the NP formalism, which is based on complex null tetrads $\{\mathbf{e}_a\} = (\mathbf{m}, \bar{\mathbf{m}}, \mathbf{n}, \bar{\mathbf{n}})$, i.e., the metric is given by

$$g_{ij} du^i du^j = \omega^1 \omega^2 - \omega^3 \omega^4, \quad \omega^2 = \bar{\omega}^1, \quad \omega^3 = \bar{\omega}^4, \quad \omega^4 = \bar{\omega}^3.$$

The connection components are labeled by NP spin coefficients:

$$\begin{aligned} -\omega_{14} &= \sigma\omega^1 + \rho\omega^2 + \tau\omega^3 + \kappa\omega^4; & \omega_{23} &= \mu\omega^1 + \lambda\omega^2 + \nu\omega^3 + \pi\omega^4; \\ -\frac{1}{2}(\omega_{12} + \omega_{34}) &= \beta\omega^1 + \alpha\omega^2 + \gamma\omega^3 + \epsilon\omega^4. \end{aligned}$$

The curvature scalars R_{bc}^α are given by $\Lambda, \Phi_{AB}, \Psi_C$, in the usual manner [17].

A proper CH₂ configuration requires a chain of proper inclusions $\mathfrak{g}_2 \subset \mathfrak{g}_1 \subset \mathfrak{g}_0 \subset \mathfrak{g}_{-1}$ so that $\mathfrak{g}_0$ is the symmetry algebra of $\tilde{R}_{bc}^\alpha$, and $\mathfrak{g}_1, \mathfrak{g}_2$ are, respectively, the symmetry algebras of $\tilde{\Gamma}_a^{\rho_1}, \tilde{\Gamma}_a^{\rho_2}$. If the Petrov type is I, II, or III, then one can fully fix the frame by setting $\Psi_0 = \Psi_4 = 0$ and then normalizing Ψ_1 or Ψ_3 to 1. In other words, $N_1 = N_0 = 0$, and hence, by Singer's criterion, every CH₁ manifold of Petrov type I, II, or III is a homogeneous space. Proper CH₁ Lorentzian manifolds must, necessarily, be of Petrov type D, N, or O. These will be studied in a future work.

A proper CH₂ configuration, if one exists, can potentially arise in 5 ways. Only in case 5.2, do we obtain a proper CH₂ configuration. A proper CH₃ configuration, if one exists, is limited to the first 3 cases. However, since in each case we rule out the existence of a proper CH₂ configuration, a proper CH₃ configuration does not exist either.

1) $\mathfrak{g}_0 = \mathfrak{so}(3)$. The curvature is that of a conformally flat perfect fluid.

$$\begin{aligned} \Psi_A &= \Phi_{01} = \Phi_{02} = \Phi_{12} = 0. & \Phi_{00} &= \Phi_{22} = 2\Phi_{11} \\ \tilde{\Gamma}_a^{\rho_1} &\in \text{span}\{\kappa - \bar{\pi}, \sigma - \bar{\lambda}, \rho - \bar{\mu}, \tau - \bar{\nu}, \alpha + \bar{\beta}, \epsilon_1 := \Re(\epsilon), \gamma_1 := \Re(\gamma)\}. \end{aligned}$$

Bianchi identities imply that all $\tilde{\Gamma}_a^{\rho_1} = 0$. Necessarily, $\mathfrak{g}_1 = \mathfrak{g}_0$. A proper configuration does not exist.

2) $\mathfrak{g}_0 = \mathfrak{so}(1, 2)$. The curvature constants are

$$\begin{aligned} \Psi_A &= \Phi_{01} = \Phi_{02} = \Phi_{12} = 0. & \Phi_{00} &= \Phi_{22} = -2\Phi_{11} \\ \tilde{\Gamma}_a^{\rho_1} &\in \text{span}\{\kappa + \bar{\pi}, \sigma + \bar{\lambda}, \rho + \bar{\mu}, \tau + \bar{\nu}, \alpha + \bar{\beta}, \epsilon_1, \gamma_1\}. \end{aligned}$$

Again, by Bianchi $\mathfrak{g}_1 = \mathfrak{g}_0$. There are no proper configurations.

3) $\mathfrak{g}_0$ is three-dimensional, generated by spins and null rotations. The curvature is that of an aligned null radiation field on a conformally flat background:

$$\begin{aligned} \Psi_A &= \Phi_{00} = \Phi_{01} = \Phi_{11} = \Phi_{02} = \Phi_{12} = 0. \\ \tilde{\Gamma}_a^{\rho_1} &\in \text{span}\{\kappa, \sigma, \rho, \tau, \alpha + \bar{\beta}, \gamma_1, \epsilon_1\}. \end{aligned}$$

The Bianchi identities imply $\kappa = \sigma = \rho = 0$, $\tau = 2\bar{\alpha} + 2\beta$. If $\tau = 0$, then $\mathfrak{g}_1 = \mathfrak{g}_0$. Performing a spin, as necessary, suppose, WLOG $\tau = \bar{\tau} \neq 0$. Hence, $\mathfrak{g}_1$ is generated by imaginary null rotations; $\tilde{\Gamma}_a^{\rho_2} \in \text{span}\{\alpha - \bar{\beta}, \gamma_2 := \Im(\gamma), \epsilon_2 := \Im(\epsilon), \lambda + \bar{\mu}, \nu_1, \pi_1\}$. The NP equations for a CH₁ configuration give $\beta = -\frac{1}{4}\tau$, $\alpha = \frac{3}{4}\tau$, $\pi_1 = -\tau$, $\epsilon_2 = 0$, $\Lambda = -24\tau^2$, $\lambda + \bar{\mu} = -\frac{2}{3}\gamma$. These constraints describe a proper CH₁ configuration. However, the additional NP equations for a proper CH₂ configuration, imply $\Phi_{22} = \frac{8}{9}\gamma_1^2 - 2\nu_1\tau$, $\gamma_2 = 0$. The last condition implies that $\mathfrak{g}_2 = \mathfrak{g}_1$. Therefore, a proper CH₂ configuration does not exist.

4) $\mathfrak{g}_0$ is generated by spins and boosts, $N_0 = 2$. Petrov type is D/O.

$$\Psi_0 = \Psi_1 = \Psi_3 = \Psi_4 = \Phi_{00} = \Phi_{01} = \Phi_{02} = \Phi_{12} = \Phi_{22} = 0.$$

$$\tilde{\Gamma}_a^{\rho_1} \in \text{span}\{\kappa, \sigma, \rho, \tau, \lambda, \mu, \nu, \pi\}.$$

We require $N_1 = 1$. There are 2 subcases. Neither admits a proper configuration.

4.1) $\mathfrak{g}_1$ = spins, $\kappa = \sigma = \lambda = \nu = \tau = \pi = 0$, $(\rho, \mu) \neq (0, 0)$. Bianchi identities imply $\Psi_2 = -\frac{2}{3}\Phi_{11}$, $\mu + \bar{\mu} = 0$, $\rho + \bar{\rho} = 0$. The NP equations, then imply $\rho^2 = 0$, and $\mu^2 = 0$, a contradiction.

4.2) $\mathfrak{g}_1$ = boosts, $\kappa = \sigma = \lambda = \nu = \rho = \mu = 0$, $(\tau, \pi) \neq (0, 0)$. By Bianchi, $\Psi_2 = \frac{2}{3}\Phi_{11}$, $\tau = \bar{\pi}$. NP equations imply $\pi^2 = 0$, a contradiction.

5) $\mathfrak{g}_0$ is 2-dimensional, generated by null rotations. The curvature is null radiation/vacuum with an aligned type N or conformally flat background:

$$\Psi_0 = \Psi_1 = \Psi_2 = \Psi_3 = \Phi_{00} = \Phi_{01} = \Phi_{11} = \Phi_{02} = \Phi_{12} = 0;$$

$$\tilde{\Gamma}_a^{\rho_1} \in \text{span}\{\kappa, \sigma, \rho, \tau, \alpha, \beta, \gamma, \epsilon\}.$$

To obtain a proper CH₂ configuration, we require $N_1 = 1$. WLOG, $\mathfrak{g}_1$ consists of imaginary null rotations; $\tilde{\Gamma}_a^{\rho_2} \in \text{span}\{\lambda + \bar{\mu}, \nu_1, \pi_1\}$. This requires $\kappa = \sigma = \rho = 0$, $\alpha = \beta + \tau \neq 0$. We use a real null rotation to set $\gamma_1 = 0$. There are two subcases: the singular case $\gamma_2 = 0$, and the generic case $\gamma_2 \neq 0$.

5.1) Suppose $\gamma_2 = 0$. The CH₁ NP equations give $\tau_2 = \beta_2 = 0$, $\Lambda = -24\tau^2$, $\bar{\Psi}_4 = \Psi_4$, $\pi_1 = -\tau_1$, $\gamma_2 = 0$, $\lambda = -\bar{\mu}$. However, the last 3 equations imply $\mathfrak{g}_2 = \mathfrak{g}_1$. Therefore, no proper CH₂ configurations are possible.

5.2) Suppose $\gamma_2 \neq 0$. The CH₁ NP equations and Bianchi identities give $\tau_2 = 0$, $\beta = \tau_1/2$, $\Lambda = -24\tau_1^2$, $\Psi_4 = 3\Phi_{22}$, $\pi_1 = -\tau_1$, $\lambda = -\bar{\mu} - \frac{2i}{3}\gamma_2$. We ask that $N_2 = 0$; $\Gamma_a^\lambda \in \text{span}(\lambda - \bar{\mu}, \nu_2, \pi_2)$. This requires $\gamma_2 \neq 0$. WLOG, an imaginary null rotation normalizes $\nu_2 = 0$, and a boost normalizes $\gamma_2 = \frac{3}{2}$. The CH₂ NP equations then imply $\pi_2 = 0$, $\mu = \frac{2i}{5}(4 + \Phi_{22})$.

The EDS for the resulting proper CH₂ configuration is generated by a Pfaffian system (1-forms and their exterior derivatives),

$$\omega_{(ab)} = 0, \quad \omega_{[14]} = -\tau_1 \omega^1, \quad \omega_{[34]} = -2\tau_1(\omega^1 + \omega^2), \quad \omega_{[12]} = \tau_1(\omega^1 - \omega^2) - 3i\omega^3,$$

$$\omega_{[13]} = i(\omega^1 - \omega^2) - \tau_1 \omega^4 - \frac{i}{5}(3 + 2\Phi_{22})(\omega^1 + \omega^2) - i\nu_2 \omega^3$$

and an additional 2-form,

$$d\nu_2 \wedge \omega^3 = \frac{3i}{5}(4 + \Phi_{22})(\omega^1 - \omega^2) \wedge \omega^3 - 3\nu_2 \tau_1(\omega^1 + \omega^2) \wedge \omega^3 + 3\tau_1 \omega^3 \wedge \omega^4.$$

Here $\tau_1 \neq 0$, $\Phi_{22} \neq 0$ are real constants. The total space is $\mathcal{F}^2 M \times \mathbb{R}$, with ν_2 the coordinate of the extra factor. Cartan's test yields $s_0 = 16$, $s_1 = 17$, $s_2 = 12$, $s_3 = 8$, $s_4 = 4$, which proves that the EDS is involutive. Subtracting (16,16,12,8,4), the parameter counts corresponding to a choice of local coordinates, we infer that, up to a local diffeomorphism, the general solution depends on 1 function of 1 variable. Indeed, an exact solution can be given:

$$\omega^1 = \frac{1}{2C_1} \left((1 + 3iu)dr + 3idu - e^r \left(3u + i \left(C_1^2 C_2 + \frac{9}{2}u^2 - \frac{5}{2} \right) \right) dx \right) \quad (24)$$

$$\omega^3 = e^r dx \quad (25)$$

$$\omega^4 = dv + 3(C_2 u + v)dr + \left(f(x)e^{-3r} + e^r \left(\frac{C_1^2 C_2^2}{4} - 9uv - \frac{5}{8}C_2(1 + 18u^2) \right) \right) dx \quad (26)$$

where x, r, u, v are coordinates; $f(x)$ is the functional parameter; C_1, C_2 are real constants such that $\tau_1 = C_1$, $\Phi_{22} = -4 + \frac{5}{2}C_1^2C_2$, $\nu_2 = -\frac{3}{2}C_1(3C_2u + 2v)$.

Finally, we note that, since $N_2 = 0$, it is impossible to refine the above to a proper CH_3 configuration.

Acknowledgments

The research of RM is supported in part by NSERC grant RGPIN-228057-2004.

References

- [1] Bueken P 1997 *J. Geom. Phys.* **22**N 349–362
- [2] Bueken P and Vanhecke L 1997 *Class. Quantum Grav.* **14** L93–L96
- [3] Bueken P and Djorić M 2000 *Ann. Global Anal. Geom.* **18** 85–103
- [4] Dolan P and Gerber A 2004 *J. Math. Phys.* **45** 310–326
- [5] Dunn C, Gilkey P and Nikčević S *Preprint* math.DG/0408316.
- [6] Estabrook F B 1987 *Acta Appl. Math.* **8** 293–319
- [7] Estabrook F B and Wahlquist H D 1989 *Class. Quantum Grav.* **6** 263–274
- [8] Estabrook F B 2006 *Class. Quantum Grav.* **23** 2841–2848
- [9] Fels M E and Renner A G 2006 *Canad. J. Math.* **58** 282–311
- [10] Gilkey P and Nikčević S 2005 *Ann. Global Anal. Geom.* **27** 87–100
- [11] Gilkey P The spectral geometry of the riemannian curvature tensor *Preprint*
- [12] Ivey T A and Landsberg J M 2003 *Cartan for beginners: differential geometry via moving frames and exterior differential systems* (Providence, RI: AMS)
- [13] Kobayashi S 1972 *Transformation groups in differential geometry* (New York, NY:Springer-Verlag)
- [14] Kowalski O, Tricerri F and Vanhecke L 1992 *J. Math. Pures Appl.*, **71** 471–501
- [15] Kowalski O and Vlášek Z 1998 *Bull. Belg. Math. Soc.* **5** 59–68
- [16] Podesta F and Spiro A Introduzione ai gruppi di trasformazioni *Preprint*
- [17] Stephani H, Kramer D, MacCallum M, Hoenselaers C and Herlt E 2003 *Exact solutions of Einstein's field equations* (Cambridge: Cambridge University Press)
- [18] Sekigawa K, Suga H and Vanhecke L 1992 *Comment. Math. Univ. Carolin.* **33** 261–268
- [19] Singer I M 1960 *Comm. Pure Appl. Math.* **13** 685–697.